arXiv:1610.05630v1 [math.AG] 18 Oct 2016

ON THE CHARACTERIZATION OF ABELIAN VARIETIES FOR LOG PAIRS IN ZERO AND POSITIVE CHARACTERISTIC

YUAN WANG

Abstract. Let (X, Δ) be a pair. We study how the condition $\kappa(K_X + \Delta) = 0$ causes surjectivity or birationality of the Albanese map and the Albanese morphism of X in both characteristic 0 and characteristic $p > 0$. In particular in characteristic 0 we generalize Kawamata's result to the cases of log canonial pairs, and in characteristic $p > 0$ we generalize a result of Hacon-Patakfalvi to the cases of log pairs. Moreover we show that if X is a normal projective threefold in characteristic $p > 0$, the coefficients of the components of Δ are ≤ 1 and $-(K_X + \Delta)$ is semiample, then the Albanese morphism of X is surjective under reasonable assumptions on p and the singularities of the general fibers of the Albanese morphism. This is a positive characteristic analog in dimension 3 of a result of Zhang on a conjecture of Demailly-Peternell-Schneider.

1. Introduction

Over the past few decades there have been many interesting discoveries on the birational geometry of the Albanese morphism in both zero and positive characteristic, especially in higher dimensions. The celebrated result [Kaw81, Theorem 1] of Kawamata shows that for a smooth complex projective variety X, if the Kodaira dimension of X is 0 then the Albanese morphism of X is an algebraic fiber space. Later Hacon and Patakfalvi established some positive characteristic analogs of Kawamata's result in [HP13] for smooth varieties and in [HP15] for possibly singular varieties of maximal Albanese dimension. It is then natural to ask whether such results hold for log pairs as well:

Question 1.1. For a pair (X, Δ) (possibly with some assumption on the singularities), does the Kodaira dimension of $K_X + \Delta$ being 0 imply that the Albanese map or the Albanese morphism of X is surjective, and do they have connected fibers?

Note that the Albanese map and the Albanese morphism can be different in general for singular varieties (see Section 2).

Another motivation originates from the following conjecture of Demailly-Peternell-Schneider:

Conjecture 1.2. [DPS93, Conjecture 2] *Let X be a compact Kähler manifold with $-K_X$ nef. Then the Albanese morphism of X is surjective.*

2010 *Mathematics Subject Classification.* 14E15, 14J10, 14J17, 14J30.

Key words and phrases. abelian variety, Albanese map, Albanese morphism, canonical bundle formula, generic vanishing, Kodaira dimension, positive characteristic.

The author was supported in part by the FRG grant DMS-#1265261, the NSF research grant DMS-#1300750 and the Simons Foundation Award # 256202.

In [Zha96], Zhang confirms Conjecture 1.2 when X is a smooth projective variety. Moreover he shows that in this case the Albanese morphism of X has connected fibers. Later Zhang generalized this result to the cases of log canonical pairs ([Zha05, Corollary 2]). More specifically he showed that if X is a projective variety and Δ is an effective $\mathbb{Q}$-divisor such that the pair (X,Δ) is log canonical and $-(K_X+\Delta)$ is nef, then the Albanese morphism from any smooth model of X is an algebraic fiber space.
From the characteristic p perspective the above results by Zhang induces the following natural question:

Question 1.3. Does a statement similar to [Zha05, Corollary 2] hold in positive characteristic?

In this paper, we answer a large part of Question 1.1 in all dimensions and answer Question 1.3 in dimension 3. In characteristic p, [HP13, Theorem 1.1.1 (1)] showed that for a smooth projective variety X, if the Frobenius stable Kodaira dimension $\kappa_S(X)$ is 0, then the Albanese morphism of X is surjective. Our first result generalizes [HP13, Theorem 1.1.1 (1)] to the cases of log pairs as follows.

Theorem A (Theorem 3.1)**.** *Let X be a normal $\mathbb{Q}$-factorial projective variety over an algebraically closed field k of characteristic $p>0$, and Δ an effective $\mathbb{Q}$-divisor on X. Assume that $\kappa_S(X+\Delta)=0$ and the Cartier index of $K_X+\Delta$ is not divisible by p. Then the Albanese morphism of X is surjective.*

The proof of Theorem A is similar to that of [HP13, Theorem 1.1.1 (1)]. Nevertheless Theorem A is interesting in the sense that to the best of the author's knowledge, there is no result of this kind in characteristic 0. We next present our positive characteristic analog of [Zha05, Corollary 2] in dimension 3 as follows. The proof of this result is inspired by some ideas in the previous paper [Wan15b] of the author.

Theorem B (Theorem 3.4)**.** *Let X be a normal projective threefold over an algebraically closed field k of characteristic $p>0$, and (X,Δ) a pair such that the coefficients of the components of Δ are ≤ 1 and $-(K_X+\Delta)$ is semiample. Suppose that*

(1) *$p>\max\{\frac{2}{\delta}, i_{\mathrm{bpf}}(-(K_X+\Delta))\}$, where δ is the minimal non-zero coefficient of Δ.*
(2) *There is a fiber X_0 of the Albanese morphism a_X of X such that $(X_0,\Delta|_{X_0})$ is normal and sharply F-pure.*

Then a_X is surjective.

Here $i_{\mathrm{bpf}}(-(K_X+\Delta))$ is the *base-point-free index* of $-(K_X+\Delta)$, which is the smallest integer $m>0$ such that $-m(K_X+\Delta)$ is base point free (cf. Definition 3.3). In particular if we assume that $\Delta=0$ and $|-K_X|$ is base point free, then condition (1) is automatically satisfied, and in that case Theorem B works in small characteristics as well.

It is worth noting that [Zha05, Corollary 2] does not hold in general if (X,Δ) is not log canonical, even in dimension 2, as is shown in Example 3.2. The example is a projective cone S over a non-hyperelliptic curve of genus ≥ 2 such that $K_S\sim_{\mathrm{lin}}0$, so obviously its Albanese map of S is not surjective (but the Albanese morphism of S is). Observe that in this case the vertex of S is its non-lc center, and the exceptional divisor over the vertex dominates the image of the Albanese map of S, which is the image of the curve in its Jacobian. This phenomenon leads us to the following result in characteristic 0.

Theorem C (Theorem 4.2)**.** *Let X be a normal complex projective variety, (X,Δ) a pair such that $\kappa(K_X+\Delta)=0$. Suppose that the Albanese map $\mathfrak{a}_X : X \dashrightarrow \mathfrak{Alb}(X)$ (respectively the Albanese morphism $a_X : X \to \mathrm{Alb}(X)$) of X is not an algebraic fiber space. Then the non-lc locus of (X,Δ) dominates $\mathfrak{a}_X(X)$ (respectively $a_X(X)$).*

The proof of Theorem C has various kinds of ingrediendts, including analyzing in detail the structure of the Albanese map and the Albanese morphism, using Campana's work on the Iitaka conjecture ([Cam04]) over a variety of general type, and applying a generic vanishing result developed by Popa and Schnell ([PS14]) to compare certain Kodaira dimensions. A direct consequence of Theorem C is that if (X,Δ) is lc and $\kappa(K_X+\Delta)=0$, then both the Albanese map and the Albanese morphism of X are algebraic fiber spaces. Moreover if the dimension of X is equal to that of $\mathfrak{Alb}(X)$ or $\mathrm{Alb}(X)$, then X is birational to an abelian variety. Therefore Theorem C generalizes [Kaw81, Theorem 1] to a great extent.

The author suspects that Theorem C is already optimal for Albanese maps, but not for Albanese morphisms, as is illustrated by Example 3.2.

Acknowledgements. This project started from a question the author put forward in Christopher Hacon's class. The author would like to thank Christopher for his constant support, encouragement and many enlightening discussions. He is also indebted to Karl Schwede who provided lots of help on F-singularities, and Sofia Tirabassi who pointed out the very important reference [Shi15]. Moreover he thanks Omprokash Das, Tommaso de Fernex, Yoshinori Gongyo, Zsolt Patakfalvi and Shunsuke Takagi for helpful conversations. Part of this project was done when the author was attending the conference "Higher Dimensional Algebraic Geometry and Characteristic p" at the Centre International de Rencontres Mathématiques and was visiting the University of Oslo, and he wishes to thank the organizers, especially John Christian Ottem, for their support and hospitality.

2. Notations and preliminaries

Throughout the article we work over an algebraically closed field k. As we will consider the cases in both characteristic 0 and characteristic $p>0$ we do not make any restriction on the characteristic of k in advance.

We set up the following conventions.

(1) For definitions related to singularities in birational geometry (e.g. *pairs*, *discrepancy*, *klt*, *lc*) we follow [HK10, Part I, Chapter 3].
(2) For a morphism $f : X \to Y$ of varieties we use both $F_{X/Y}$ and F_f to denote a general fiber of f, and we say that f is an *algebraic fiber space* if it is surjective and has connected fibers.

2.1. Albanese variety, Albanese map, Albanese morphism, derived categories, Fourier-Mukai transform and generic vanishing.

We first present the definitions for the Albanese map and the Albanese moprhism. These can be found in [Lan83] and [FGI+05].

Theorem-Definition 2.1. Let X be a variety over k. There exists an abelian variety $\mathfrak{Alb}(X)$ together with a rational map $\mathfrak{a}_X : X \dashrightarrow \mathfrak{Alb}(X)$ such that

- $\mathfrak{a}_X(X)$ generates $\mathfrak{Alb}(X)$, i.e. $\mathfrak{a}_X(X)$ is not contained in a translate of any proper abelian subvariety of $\mathfrak{Alb}(X)$.
- For every rational map $f : X \dashrightarrow A$ from X to an abelian variety A, there exists a homomorphism $g : \mathfrak{Alb}(X) \to A$ and a constant $a \in A$ such that $f = g \circ \mathfrak{a}_X + a$.

We call $\mathfrak{a}_X$ the *Albanese map* of X and $\mathfrak{Alb}(X)$ the *Albanese variety via the Albanese map (AVAMa)* of X.

Theorem-Definition 2.2. Let X be a normal projective variety over k. There exists an abelian variety $\mathrm{Alb}(X)$ together with a morphism $a_X : X \to \mathrm{Alb}(X)$ such that

- $a_X(X)$ generates $\mathrm{Alb}(X)$, i.e. $a_X(X)$ is not contained in a translate of any proper abelian subvariety of $\mathrm{Alb}(X)$.
- For every morphism $f : X \to A$ from X to an abelian variety A, there exists a homomorphism $h : \mathrm{Alb}(X) \to A$ and a constant $b \in A$ such that $f = h \circ a_X + b$.

We call $\mathfrak{a}_X$ the *Albanese morphism* of X and $\mathfrak{Alb}(X)$ the *Albanese variety via the Albanese morphism (AVAMo)* of X.

Throughout the article, unless otherwise stated, for a normal projective variety X we use $\mathfrak{a}_X : X \dashrightarrow \mathfrak{Alb}(X)$ ($a_X : X \to \mathrm{Alb}(X)$) to denote the Albanese map (Albanese moprhism) of X. We refer to [Lan83, Chapter II, §3] and [FGI+05, Chapter 9] for more details about the Albanese map and the Albanese morphism respectively. The Albanese map and the Albanese morphism agree in characteristic 0 for normal proper varieties with rational singularities (cf. [Rei83, Proposition 2.3] or [Kaw85, Lemma 8.1]). But they do not agree in general, as is illustrated by the following

Example 2.3. Let X be a projective cone over any curve C of genus ≥ 1. Since X is covered by rational curves passing through the vertex and the Albanese morphism contracts all the rational curves, we see that the Albanese morphism of X has to be a morphism from X to a point. On the other hand let X' be the blow-up of X at the vertex, $\nu : X \dashrightarrow X'$ the natural birational map, $p : X' \to C$ the natural $\mathbb{P}^1$ fibration from X' to C and $j : C \hookrightarrow \mathrm{Jac(C)}$ the embedding from C to its Jacobian. Then the Albanese map of X is $j \circ p \circ \nu$, whereas the Albanese morphism (and also the Albanese map) of X' is $j \circ p$.

Example 2.3 is also an example of the fact that the AVAMa is a birational invariant, but the AVAMo is not. Still, by definition for a normal projective variety X there exists a surjective homomorphism α from $\mathfrak{Alb}(X)$ to $\mathrm{Alb}(X)$ such that $a_X = \alpha \circ \mathfrak{a}_X$.

For convenience, in characteristic 0 we say that *the Albanese map of X is an algebraic fiber space* if the Albanese morphism of any smooth model of X is an algebraic fiber space.

Definition 2.4. For an abelian variety A and a coherent sheaf $\mathcal{F}$ on A, we define $V^i(\mathcal{F})$, the *i-th cohomology support locus*, as

$$V^i(\mathcal{F}) = \{P \in \mathrm{Pic}^0(A) | h^i(A, \mathcal{F} \otimes P) > 0\}.$$

$\mathcal{F}$ is called a *GV-sheaf*, if $\mathrm{codim}_{\mathrm{Pic}^0(A)} V^i(\mathcal{F}) \geq i$ for all $i \geq 0$.

Theorem 2.5. [Hac04, Corollary 3.2 (1)] *Let $\mathcal{F}$ be a GV-sheaf on an abelian variety A. For any $P \in \hat{A}$, if $i \geq 0$ and $H^i(A, \mathcal{F} \otimes P) = 0$, then $H^{i+1}(A, \mathcal{F} \otimes P) = 0$.*

Theorem 2.6. [PS14, Variant 5.5] *Let $f : X \to A$ be a morphism from a normal projective variety to an abelian variety. If (X, Δ) is a lc pair and $k > 0$ is any integer such that $k(K_X + \Delta)$ is Cartier, then $f_*\mathcal{O}_X(k(K_X + \Delta))$ is a GV-sheaf.*

For a normal projective variety of dimension n, we denote by $D(X)$ the derived category of $\mathcal{O}_X$ modules, and denote by $D_{\mathrm{qc}}(X)$ the full subcategory of $D(X)$ consisting of bounded complexes with quasi-coherent cohomologies. We denote the dualizing complex of X by $\omega_X^{\cdot}$ and define the dualizing functor D_X by $D_X(F) \coloneqq R\mathcal{H}om(F, \omega_X^{\cdot}[n])$ for any $F \in D_{\mathrm{qc}}(X)$.

Definition 2.7. [Muk81] Let A be an abelian variety and $\hat{A}$ its dual abelian variety. Let $p_A : A \times \hat{A} \to A$ and $p_{\hat{A}} : A \times \hat{A} \to \hat{A}$ be the natural projection morphisms and $\mathcal{P}$ the Poincaré line bundle on $A \times \hat{A}$. We define the *Fourier-Mukai transform* $R\hat{S} : D_{\mathrm{qc}}(A) \to D_{\mathrm{qc}}(\hat{A})$ and $RS : D_{\mathrm{qc}}(\hat{A}) \to D_{\mathrm{qc}}(A)$ with respect to the kernel $\mathcal{P}$ by

$$R\hat{S}(\cdot) = Rp_{\hat{A},*}(p_A^*(\cdot) \otimes \mathcal{P}), \quad RS(\cdot) = Rp_{A,*}(p_{\hat{A}}^*(\cdot) \otimes \mathcal{P}).$$

For an abelian variety A of dimension g, $R^g\hat{S}$ gives an equivalence in categories between unipotent vector bundles on A and the category of Artinian $\mathcal{O}_{\hat{A},\hat{0}}$-modules of finite length on $\hat{A}$ (cf. [CH09, Section 1.2]).

Lemma 2.8. *Suppose that* $\mathrm{char}(k) = 0$. *Let (X, Δ) be a log canonical pair and $f : X \to A$ an algebraic fiber space from X to an abelian variety A. If $\kappa(K_{F_f} + \Delta|_{F_f}) \geq 0$, then $\kappa(K_X + \Delta) \geq 0$.*

Proof. If $\kappa(K_{F_f} + \Delta|_{F_f}) \geq 0$, then $H^0(F_f, \mathcal{O}_{F_f}(m(K_{F_f} + \Delta|_{F_f}))) \neq 0$ for a sufficiently divisible m such that $m(K_X + \Delta)$ is Cartier. So by cohomology and base change we have that $f_*(\mathcal{O}_X(m(K_X + \Delta))) \neq 0$.

Next we prove that $V^0(f_*\mathcal{O}_X(m(K_X + \Delta))) \neq 0$. By Theorem 2.6 we know that $f_*\mathcal{O}_X(m(K_X + \Delta))$ is a GV sheaf, in particular

$$V^i(f_*\mathcal{O}_X(m(K_X + \Delta))) \supseteq V^{i+1}(f_*\mathcal{O}_X(m(K_X + \Delta)))$$

for all $i \geq 0$. Suppose that $V^0(f_*\mathcal{O}_X(m(K_X + \Delta))) = 0$. Then $V^i(f_*\mathcal{O}_X(m(K_X + \Delta))) = 0$ for all i. By cohomology and base change we have $R\hat{S}(f_*\mathcal{O}_X(m(K_X + \Delta))) = 0$ (cf. [Wan15a, Proof of Proposition 2.13]). So by [Muk81, Theorem 2.2] we have $f_*(\mathcal{O}_X(m(K_X + \Delta))) = 0$, which contradicts what we deduced in the last paragraph.

So we have that $V^0(f_*\mathcal{O}_X(m(K_X + \Delta))) \neq 0$. On the other hand, by [Shi15, Theorem 1.3], $V^0(f_*\mathcal{O}_X(m(K_X + \Delta)))$ is a finite union of torsion translates of abelian subvarieties of $\mathrm{Pic}^0(A)$. In particular we have

$$0 \neq H^0(A, f_*\mathcal{O}_X(m(K_X + \Delta)) \otimes P) = H^0(X, \mathcal{O}_X(m(K_X + \Delta)) \otimes f^*P)$$

for some torsion elements $P \in \mathrm{Pic}^0(A)$. This implies that $H^0(X, \mathcal{O}_X(m'(K_X + \Delta))) \neq 0$ for sufficiently divisible m', hence $\kappa(K_X + \Delta) \geq 0$. □

2.2. **F-singularities.** We recall the definitions of some basic concepts of F-singularity. We refer to [ST11], [PST14], [HX15] and [DS15] for more details. Let k be of characteristic $p > 0$ and X a normal projective variety over k. Let (X, Δ) be a pair such that the Cartier index of $K_X + \Delta$ is not divisible by p. Let

$$\mathcal{L}_{e,\Delta} \coloneqq \mathcal{O}_X((1 - p^e)(K_X + \Delta))$$

where e is a positive integer such that $(p^e-1)(K_X+\Delta)$ is Cartier. Then there is a canonically determined morphism $\phi_e : F_*^e\mathcal{L}_{e,\Delta} \to \mathcal{O}_X$ (cf. [PST14, Proposition 3.4]). We define the *non-F-pure ideal* $\sigma(X,\Delta)$ (the *test ideal* $\tau(X,\Delta)$) to be the unique biggest (smallest) ideal $J \subseteq \mathcal{O}_X$ such that $(\phi_e \circ F_X^e)(J\cdot\mathcal{L}_{e,\Delta}) = J$ for any e. We say that (X,Δ) is *sharply F-pure* (*strongly F-regular*) if $\sigma(X,\Delta) = \mathcal{O}_X$ ($\tau(X,\Delta) = \mathcal{O}_X$). For an integral subscheme $Z \subseteq X$ with generic point η_Z, we say that Z is an *F-pure center* of (X,Δ) if

- Δ is effective at η_Z,
- (X,Δ) is F-pure at η_Z, and
- $\phi(F_*^e I_Z\cdot\mathcal{L}_{e,\Delta})\cdot\mathcal{O}_{X,\eta_Z} \subseteq I_Z\cdot\mathcal{O}_{X,\eta_Z}$.

Lemma 2.9. *Let (X,Δ) be a sharply F-pure pair such that the Cartier index of $(K_X+\Delta)$ is not divisible by p. Let D be a divisor such that D does not contain any F-pure center of (X,Δ). Then there is a positive integer n such that $(X,\Delta+\frac{1}{n}D)$ is sharply F-pure.*

Proof. We first construct a stratification for the F-pure centers of (X,Δ). Let $W_i^*(X,\Delta)$ be the union of $\le i$-dimensional F-pure centers of (X,Δ). Let

$$W_i(X,\Delta) := W_i^*(X,\Delta)\setminus W_{i-1}^*(X,\Delta)$$

and $W_i^k(X,\Delta)$ the irreducible components of $W_i(X,\Delta)$. Then X is a disjoint union of $W_i^k(X,\Delta)$, and each $W_i^k(X,\Delta)$ is a minimal F-pure center of (X,Δ). In particular $W_i^k(X,\Delta)$ is normal. For each $W_i^k(X,\Delta)$, by [Sch09, (i) and (iv) of Main Theorem] there is an effective divisor $\Delta_{W_i^k(X,\Delta)}$ on $W_i^k(X,\Delta)$ such that $(K_X+\Delta)|_{W_i^k(X,\Delta)} = K_{W_i^k(X,\Delta)} + \Delta_{W_i^k(X,\Delta)}$ and $(W_i^k(X,\Delta),\Delta_{W_i^k(X,\Delta)})$ is strongly F-regular. Then there is a positive integer $n_i^k > 0$ such that $(W_i^k(X,\Delta),\Delta_{W_i^k(X,\Delta)}+\frac{1}{n_i^k}D|_{W_i^k(X,\Delta)})$ is strongly F-regular. Hence by [Sch09, (iii) of Main Theorem], $(X,\Delta+\frac{1}{n_i^k}D)$ is sharply F-pure near $W_i^k(X,\Delta)$. Finally let $n := \max\{n_i^k\}$ and we are done. □

Let $f : X \to Y$ be a morphism between normal projective varieties and $\mathcal{M}$ a line bundle on X. We define the subsheaf $S^0f_*(\sigma(X,\Delta)\otimes\mathcal{M}) \subseteq f_*\mathcal{M}$ as

$$S^0f_*(\sigma(X,\Delta)\otimes\mathcal{M}) := \bigcap_{g\ge 0}\mathrm{Image}(F_*^{ge}f_*(\mathcal{L}_{ge,\Delta}\otimes\mathcal{M}^{p^{ge}}) \xrightarrow{\Phi_{ge}} f_*\mathcal{M}) \tag{2.1}$$

where Φ_{ge} is induced by ϕ_{ge} using the projection formula and the commutativity of F and f. In the case $Y = \mathrm{Spec}\ k$, $f_*(\cdot)$ is the same as $H^0(\cdot)$, and in this case we use $S^0(X,\Delta;\mathcal{M})$ to denote $S^0f_*(\sigma(X,\Delta)\otimes\mathcal{M})$. Since $H^0(X,\mathcal{M})$ is finite dimensional, we have

$$S^0(X,\Delta;\mathcal{M}) = \mathrm{Image}(H^0(X,F_*^{me}(\mathcal{L}_{me,\Delta}\otimes\mathcal{M}^{p^{me}})) \xrightarrow{\Phi_{me}} H^0(X,\mathcal{M}))$$

for some $m \gg 0$.

The following definition is analogous to that in [HP13, 4.1].

Definition 2.10. Let (X,Δ) be a pair and $r > 0$ an integer such that $r(K_X+\Delta)$ is Cartier. The *Frobenius stable Kodaira dimension* of (X,Δ) is defined as

$$\begin{aligned}&\kappa_S(K_X+\Delta)\\ &:= \max\{k \mid \dim S^0(X,\Delta;\mathcal{O}_X(mr(K_X+\Delta))) = O(m^k) \text{ for } m \text{ sufficiently divisible}\}.\end{aligned}$$

It is easy to see that $\kappa_S(K_X+\Delta)$ is independent of r.

2.3. **A canonical bundle formula in characteristic p.** The following theorem is a slight modification of [Wan15b, Theorem 4.3], which is a canonical bundle formula for morphism from a threefold to a surface whose general fibers are $\mathbb{P}^1$. The same proof in [Wan15b] works.

Theorem 2.11. *Let $f: X \to Y$ be a proper surjective morphism, where X is a normal threefold and Y is a normal surface over an algebraically closed field k of characteristic $p>0$. Assume that $Q=\sum_i Q_i$ is a divisor on Y such that f is smooth over $(Y-\mathrm{Supp}(Q))$ with fibers isomorphic to $\mathbb{P}^1$. Let D be an effective $\mathbb{Q}$-divisor on X, which satisfies the following conditions:*

1. *(X,D) is lc on a general fiber of f.*
2. *Suppose $D=D^h+D^v$ where D^h is the horizontal part and D^v is the vertical part of D. Then $p=\mathrm{char}(k)>\dfrac{2}{\delta}$, where δ is the minimum non-zero coefficient of D^h.*
3. *$K_X+D\sim_{\mathbb{Q}} f^*(K_Y+M)$ for some $\mathbb{Q}$-Cartier divisor M on Y.*

Then M is $\mathbb{Q}$-linearly equivalent to an effective $\mathbb{Q}$-divisor.

3. Surjectivity of the Albanese maps in characteristic $p>0$

In this section we first prove the following theorem, which generalizes [HP13, Theorem 1.1.1 (1)] to the cases of log pairs.

Theorem 3.1. *Let X be a normal $\mathbb{Q}$-factorial projective variety in characteristic $p>0$, and Δ an effective $\mathbb{Q}$-divisor on X. Assume that $\kappa_S(K_X+\Delta)=0$ and the Cartier index of $K_X+\Delta$ is not divisible by p. Then the Albanese morphism of X is surjective.*

Proof. For convenience, in this proof we use $a: X\to A$ to denote the Albanese morphism of X. $\kappa_S(K_X+\Delta)=0$ implies that there is a positive integer r such that $r(K_X+\Delta)$ is Cartier and $S^0(X,\Delta;\mathcal{O}_X(r(K_X+\Delta)))\neq 0$. We fix an $e>0$ such that $r|(p^e-1)$. Then for an $m\gg 0$ there is an element of $H^0(X,\mathcal{O}_X((p^{me}r+(1-p^{me}))(K_X+\Delta)))$ mapping to $0\neq f\in H^0(X,\mathcal{O}_X(r(K_X+\Delta)))$. By [HP13, Lemma 4.1.3] we know that $\kappa(K_X+\Delta)=0$, so that element can only be $\alpha f^{p^{me}+\frac{1-p^{me}}{r}}$ for some $\alpha\in k^*$. Let $G\in|r(K_X+\Delta)|$ be the unique element and $D:=\frac{r-1}{r}G$. Since the image of

$$\alpha f\in H^0(X,\mathcal{O}_X(r(K_X+\Delta)))=H^0(X,\mathcal{O}_X(rp^{me}(K_X+\Delta)+(1-p^{me})(K_X+\Delta+D)))$$

in $H^0(X,\mathcal{O}_X(r(K_X+\Delta)))$ is computed by

$$\alpha f\mapsto \alpha f\cdot f^{\frac{r-1}{r}(p^{me}-1)}=\alpha f^{p^{me}+\frac{1-p^{me}}{r}}\mapsto \beta f,$$

we have actually proved that $S^0(X,\Delta+D;\mathcal{O}_X(r(K_X+\Delta)))\neq 0$. Then the natural inclusion (cf. [HP13, Lemma 2.2.2])

$$S^0(X,\Delta+D;\mathcal{O}_X(r(K_X+\Delta)))\subseteq H^0(A,\Omega_0)\subseteq H^0(X,\mathcal{O}_X(r(K_X+\Delta)))$$

are equalities, where $\Omega_0=S^0a_*(\sigma(X,\Delta+D)\otimes\mathcal{O}_X(r(K_X+\Delta)))$. We consider the following diagram

$$\begin{array}{ccc}
F_*^{(m+1)e}\Omega_0 & \longrightarrow & F_*^{me}\Omega_0 \\
\downarrow & & \downarrow \\
F_*^{(m+1)e}a_*\mathcal{O}_X(r(K_X+\Delta)) & \longrightarrow & F_*^{me}a_*\mathcal{O}_X(r(K_X+\Delta)) \\
\downarrow{\scriptstyle =} & & \downarrow{\scriptstyle =} \\
a_*F_*^{(m+1)e}\mathcal{O}_X(r(K_X+\Delta)) & \longrightarrow & a_*F_*^{me}\mathcal{O}_X(r(K_X+\Delta))
\end{array}$$

where the bottom two horizontal arrows are from [Pat13, Lemma 2.6]. After applying $H^0(A,-)$ we get that the bottom row is an isomorphism as the stable image of these maps is exactly $S^0(X,\Delta+D;\mathcal{O}_X(r(K_X+\Delta))) = H^0(X,\mathcal{O}_X(r(K_X+\Delta)))$, which is proved above. Therefore the natural maps

$$H^0(A,F_*^{(m+1)e}\Omega_0) \to H^0(A,F_*^{me}\Omega_0)$$

and

$$H^0(X,F_*^{(m+1)e}\mathcal{O}_X(r(K_X+\Delta))) \to H^0(X,F_*^{me}\mathcal{O}_X(r(K_X+\Delta)))$$

are isomorphisms.

Now by assumption there is a unique $G \in |r(K_X+\Delta)|$. We use the following notation

$$\Omega_m := F_*^{me}S^0a_*(\sigma(X,\Delta+D)\otimes\mathcal{O}_X(r(K_X+\Delta))) \text{ and } \Omega = \varprojlim \Omega_m.$$

It is easy to see that $r|(p^{me}-1)$ for every $m \geq 0$. According to [Pat13, Lemma 2.6] and [BS13, Lemma 8.1.4] we know that $\Omega_{(m+1)} \to \Omega_m$ is surjective for every $m \geq 0$. Further by the above argument, $S^0a_*(\sigma(X,\Delta+D)\otimes\mathcal{O}_X(r(K_X+\Delta))) \neq 0$. So for every m, $\Omega_m \neq 0$, hence $\Omega \neq 0$.

Next we claim that since $\kappa(K_X+\Delta)=0$, there is a neighborhood U of the origin of $\hat{A}$ such that

$$V^0(\mathcal{O}_X(r(K_X+\Delta))) \cap U = 0_{\hat{A}}.$$

Let $g := \dim A$. Suppose that this is not the case and let $T \in \hat{A}$ be a positive dimensional irreducible component of $V^0(\mathcal{O}_X(r(K_X+\Delta)))$ through the origin. Let $\xi : T^{g+1} \to \hat{A}$ be the natural morphism. For dimensional reason, every fiber of ξ is of positive dimension. Moreover since $0 \in T$, 0 is in the image of ξ. Consider for $(P_1,...,P_{g+1}) \in \xi^{-1}(\mathcal{O}_A)$ the maps

$$\begin{aligned}
&H^0(X,\mathcal{O}_X(r(K_X+\Delta))\otimes a^*P_1)\otimes ...\otimes H^0(X,\mathcal{O}_X(r(K_X+\Delta))\otimes a^*P_{g+1}) \\
&\to H^0(X,\mathcal{O}_X((g+1)r(K_X+\Delta))).
\end{aligned}$$

Since $h^0(X,\mathcal{O}_X((g+1)r(K_X+\Delta))) = 1$, there are only finitely many choices for the divisors in $|r(K_X+\Delta)+a^*P_i|$ for $P_i \in p_i(\xi^{-1}(\mathcal{O}_A))$. However there exists an i such that $p_i(\xi^{-1}(\mathcal{O}_A))$ is an infinite set, and we also have $(\mathrm{Pic}^0(X))_{\mathrm{red}} = \mathrm{Pic}^0(A) = \hat{A}$ (cf. [Moc12, Proposition A.6]). In particular the map $\mathrm{Pic}^0(A) \to \mathrm{Pic}^0(\mathrm{X})$ is injective. So we get a contradiction, hence the claim holds.

Since $H^0(A,S^0a_{X,*}(\sigma(X,\Delta+D)\otimes\mathcal{O}_X(r(K_X+\Delta)))\otimes P) \subseteq H^0(X,\mathcal{O}_X(r(K_X+\Delta))\otimes a^*P)$ we have that $H^0(A,\Omega_0\otimes P)$ is zero for every $\mathcal{O}_A \neq P \in U$. Moreover $H^0(A,\Omega_0) \neq 0$ as shown

above, so by [HP13, Corollary 3.2.1] we have that $\mathcal{H}^0(\Lambda_0)$ is Artinian in a neighborhood of the origin and $\mathcal{H}^0(\Lambda_0)\otimes k(0)\neq 0$, where

$$\Lambda_m := R\hat{S}(D_A(\Omega_m)).$$

Therefore we can write $\mathcal{H}^0(\Lambda_0)$ as $\mathcal{H}^0(\Lambda_0)=\mathcal{C}_0\oplus\mathcal{B}_0$, where $\mathcal{B}_0$ is Artinian and supported at 0 and $0\notin\operatorname{Supp}\mathcal{C}_0$. This induces a similar decomposition $\mathcal{C}_m\oplus\mathcal{B}_m$ of $\mathcal{H}^0(\Lambda_m)\cong V^*\mathcal{H}^0(\Lambda_0)$, where V is the Verschiebung isogeny (cf. [MvdG, (5.18) Definition]). Furthermore, the morphism $\mathcal{H}^0(\Lambda_m)\to\mathcal{H}^0(\Lambda_{m+1})$ is the direct sum of morphisms $\mathcal{C}_m\to\mathcal{C}_{m+1}$ and $\mathcal{B}_m\to\mathcal{B}_{m+1}$. Hence

$$\Lambda := \varinjlim\Lambda_m = (\varinjlim\mathcal{C}_m)\oplus(\varinjlim\mathcal{B}_m),$$

where $\mathcal{B}_m$ is Artinian for every $m\geq 0$.

Now by [HP13, Corollary 3.2.1] we have $\mathcal{H}_m\otimes k(0)\cong\mathcal{B}_m\otimes k(0)\cong H^0(A,F_*^m\Omega_0)^\vee$, so the morphism $\mathcal{B}_m\otimes k(0)\to\mathcal{B}_{m+1}\otimes k(0)$ can be identified with $H^0(A,F_*^m\Omega_0)^\vee\to H^0(A,F_*^{m+1}\Omega_0)^\vee$. However by the above argument the latter morphism is an isomorphism, and $H^0(A,\Omega_0)\neq 0$. Therefore we know that $\varinjlim(\mathcal{B}_m\otimes k(0))$, hence $\varinjlim\mathcal{B}_m$, is nonzero. Finally by [HP13, Corollary 3.2.3] we are done. □

The second main theorem in this section is a characteristic p analog of [Zha05, Corollary 2] in dimension 3. Before we present that we would like to point out that [Zha05, Corollary 2] does not hold without the assumption that (X,D) is lc. This is illustrated by the following

Example 3.2. Let C be a smooth nonhyperelliptic curve of genus $g\geq 2$. By [Har77, Proposition 5.2], K_C is very ample. Let $C\hookrightarrow\mathbb{P}^r$ be the embedding induced by $|K_C|$. Let S be the projective cone over C, $\nu: S'\to S$ the blow-up at the vertex of S, $\pi: S'\to C$ the induced morphism and E the exceptional divisor of ν. We then have that $E^2=-\deg K_C=2-2g$, and we can write

$$\nu^*K_S = K_{S'}+\lambda E$$

for some λ. By adjunction formula we see that

$$2g-2=(K_{S'}+E)\cdot E=(1-\lambda)E^2=(\lambda-1)(2g-2),$$

so $\lambda=2$. If we denote a fiber of π by f, then

$$\begin{aligned}h^0(S,\mathcal{O}_S(K_S)) &= h^0(S',\nu^*\mathcal{O}_S(K_S)) = h^0(S',\mathcal{O}_{S'}(K_{S'}+2E))\\ &= h^0(S',\mathcal{O}_{S'}(-2E+(2g-2-\deg K_C)f+2E)) = h^0(S',\mathcal{O}_{S'})=1,\end{aligned}$$

where the third equality is implied by the fact that $K_{S'}\sim_{\text{lin}}-2E+(2g-2-\deg K_C)f$ as K_C is very ample. In the above calculation we actually get that $\nu^*\mathcal{O}_S(K_S)=\mathcal{O}_S$, Hence $K_S\sim_{\text{lin}}0$. In particular $-K_S$ is nef and $\kappa(S)=0$. On the other hand by the construction of S we know that its Albanese map is not surjective. However by the argument in Example 2.3, the Albanese morphism of S is an algebraic fiber space, in particular surjective.

We now show the second main theorem. For convenience we make the following definition.

Definition 3.3. For a $\mathbb{Q}$-Cartier divisor D on a projective variety X, we define the *base-point-free index* of D to be the smallest number $m>0$ such that $|mD|$ is base point free and denote it as $i_{\text{bpf}}(D)$.

Theorem 3.4. *Let X be a normal projective threefold over an algebraically closed field k of characteristic $p > 0$, and (X, Δ) a pair such that the coefficients of the components of Δ are ≤ 1 and $-(K_X + \Delta)$ is semiample. Suppose that*

(1) *$p > \max\{\frac{2}{\delta}, i_{\mathrm{bpf}}(-(K_X + \Delta))\}$, where δ is the minimal non-zero coefficient of Δ.*
(2) *There is a fiber X_0 of the Albanese morphism a_X of X such that X_0 is normal and $(X_0, \Delta|_{X_0})$ is sharply F-pure.*

Then a_X is surjective.

Lemma 3.5. *Let X be a normal projective surface. Suppose that X has at most log canonical singularities and $\kappa(X) = 0$. Then the Albanese map (hence the Albanese morphism) of X is surjective.*

Proof. Let $\mu : X' \to X$ be a minimal resolution of singularities for X, then we have $\kappa(X') \leq \kappa(X)$.

If $\kappa(X') = 0$ then this is straight forward by the Enriques-Kodaira classification.

If $\kappa(X') = -\infty$ then there is a birational morphism $g : X' \to X''$ where X'' is a ruled surface over a curve C with a $\mathbb{P}^1$ fibration $\pi : X'' \to C$. We can write

$$K_{X'} = \pi^* K_X - \sum_i a_i E_i = g^* K_{X''} + \sum_j b_j F_j$$

where $a_i, b_j > 0$ and E_i and F_j are exceptional curves on X' for μ and g respectively. By assumption we have

$$\begin{aligned} 0 \neq H^0(X, \mathcal{O}_X(mK_X)) &= H^0(X', \mu^* \mathcal{O}_X(mK_X)) = H^0(X', \mathcal{O}_X(m(K_{X'} + \sum_i a_i E_i))) \\ &= H^0(X', \mathcal{O}_X(m(g^* K_{X''} + \sum_j b_j F_j + \sum_i a_i E_i))) \\ &\subseteq H^0(X'', \mathcal{O}_{X''}(m(K_{X''} + \sum_i a_i g_* E_i))) \end{aligned}$$

for any sufficiently divisible m. Since X'' is a ruled surface, there must be an i such that $g_* E_i$ dominates the base curve C. Since X is log canonical, by [Kol13, 3.30] the genus of E_i, hence that of C, is at most 1. In particular C is surjective onto its Jacobian. So according to the structure of $a_{X''} : X'' \to \mathrm{Alb}(X'')$ explained in Example 2.3 we are done. □

Proof of Theorem 3.4. Let $X \xrightarrow{a_X} Z = a_X(X) \subseteq \mathrm{Alb}(X)$ be the Albanese morphism of X. We denote the Stein factorization of a_X by

$$X \xrightarrow{p} Y \xrightarrow{q} Z.$$

We can assume that Y is normal. Note that by condition (2) we have that a_X, hence q, is separable. It suffices to show that the Albanese morphism of Y is surjective, and the proof of this fact is the same as that of Lemma 4.4 below. We consider the following cases according to the dimension of Y (or equivalently, that of Z).

Case 1. If $\dim Y = 0$, this is trivial.

Case 2. If $\dim Y = 1$, then Y is smooth. Since $|-m(K_X+\Delta)|$ is base point free, we can choose a suitable prime divisor $D \sim_{\text{lin}} -m(K_X+\Delta)$ such that $p > m$ and $D|_{X_0}$ does not contain any F-pure center of $(X_0, \Delta|_{X_0})$. Then by Lemma 2.9 there is a positive integer n such that $(X_0, \Delta|_{X_0} + \frac{1}{n}D|_{X_0})$ is sharply F-pure. Then by applying [PSZ13, Theorem B] on the projection $X_0 \times \mathbb{P}H^0(X_0, \mathcal{O}_{X_0}(D|_{X_0})) \to \mathbb{P}H^0(X_0, \mathcal{O}_{X_0}(D|_{X_0}))$ we know that there is a positive integer n such that $(X_0, \Delta|_{X_0} + \frac{1}{n}D_{X_0})$ is sharply F-pure for any divisor $D_{X_0} \in |D|_{X_0}|$ in a neighborhood of $D|_{X_0}$. In particular the pair

$$(X_0, \Delta|_{X_0} + \frac{1}{mn(\dim X_0+1)}(D^1_{X_0} + \ldots + D^{\dim X_0+1}_{X_0}))$$

is sharply F-pure for any $D^i_{X_0} \in |D|_{X_0}|$ in a neighborhood of $D|_{X_0}$. So by using the techniques in [Tan15, Idea of Theorem 1] we get that there is a $\mathbb{Q}$-divisor $D' \sim_{\mathbb{Q}} \frac{1}{m}D$ such that $(X_0, (\Delta+D')|_{X_0})$ is sharply F-pure. Since X_0 is normal it is S_2, G_1 and regular in codimension 1. Moreover we have $K_X + \Delta + D' \sim_{\mathbb{Q}} K_X + \Delta + \frac{1}{m}D \sim_{\mathbb{Q}} 0$. So by [Pat14, Theorem 3.10] we have that $K_{X/Y} + \Delta + D'$ is nef, hence $-K_Y$ is nef. Since Z generates $\text{Alb}(X)$, Y and Z have to be elliptic curves and we are done.

Case 3. If $\dim Y = 2$, we denote a general fiber of p by F. By assumption F is a smooth curve. We observe that $\kappa(Y) \geq 0$, otherwise Y, hence Z, is covered by rational curves and this would be a contradiction to the construction that $Z \subseteq \text{Alb}(X)$. We consider the following subcases.

Subcase 1. If $g(F) = 0$, then $F \cong \mathbb{P}^1$. In this case we can choose the prime divisor $D \sim_{\text{lin}} -m(K_X+\Delta)$ such that $(F, (\Delta + \frac{1}{m}D)|_F)$ is lc. By Theorem 2.11 there is an effective $\mathbb{Q}$-divisor M on Y such that

$$0 \sim_{\mathbb{Q}} K_X + \Delta + \frac{1}{m}D \sim_{\mathbb{Q}} p^*(K_Y + M),$$

hence $K_Y + M \sim_{\mathbb{Q}} 0$ and in particular $\kappa(Y) \leq 0$. Let $\nu : Y' \to Y$ be a minimal resolution of Y. If $M \neq 0$ or Y has non-canonical singularity, then $\kappa(Y') < 0$ and Y' is birationally ruled. This is a contradiction to the fact that Y has a finite map $q : Y \to Z \subseteq \text{Alb}(X)$ onto its image. Therefore Y has canonical singularities and by Lemma 3.5 we are done.

Subcase 2. If $g(F) \geq 1$, then after possibly doing a base change and a resolution, WLOG we can assume that both X and Y are smooth. Then by [CZ15, 3.2] we have $\kappa(\omega_{X/Y}) \geq 0$. This together with $\kappa(X) \leq 0$ implies that $\kappa(Y) \leq 0$. Again by the argument in *Subcase 1* and Lemma 3.5 we are done.

Case 4. If $\dim Y = 3$ then the a_X is generically finite. Moreover since X_0 is normal it is reduced, hence a_X is separable. So by [HP15, Theorem 0.2] we are done. □

4. Characterization of abelian varieties for lc paris in characteristic 0

In this section we work in characteristic 0.

Definition 4.1. Let (X, Δ) be a pair and $a : X \dashrightarrow Y$ a rational map from X to another variety Y . We say that *the non-lc locus of* (X, Δ) *dominates* $a(X)$ if there is a divisor E of discrepancy < -1 that dominates $a(X)$.

The following theorem is the main theorem of this section.

Theorem 4.2. *Let X be a normal complex projective variety, (X, Δ) a pair such that $\kappa(K_X + \Delta) = 0$. Suppose that the Albanese map $\mathfrak{a}_X : X \dashrightarrow \mathfrak{Alb}(X)$ (the Albanese morphism $a_X : X \to \mathrm{Alb}(X)$) of X is not an algebraic fiber space. Then the non-lc locus of (X, Δ) dominates $\mathfrak{a}_X(X)$ ($a_X(X)$).*

Lemma 4.3. *Let $f : X \to Z$ be a morphism of varieties. Let E a divisor over X and $g : Z' \to Z$ a birational morphism. Then E dominates Z if and only if E dominates Z'.*

Proof. Obvious. □

Proof of Theorem 4.2. We first deal with the case of the Albanese map. The case of Albanese moprhism is almost the same and will be explained at the end of the proof. Let $\mu : Y \to X$ be a log resolution of (X, Δ) and we can write

$$K_Y + \tilde{\Delta} = \mu^*(K_X + \Delta) + E_+ - E_-$$

where $\tilde{\Delta}$ is the strict transform of Δ and E_+ and E_- are effective exceptional divisors of μ with no common components. Denote $D \coloneqq \tilde{\Delta} + E_-$ and for convenience we denote the Albanese morphism of Y by a as well. We have $\kappa(K_Y + D) = \kappa(K_X + \Delta) = 0$ (cf. [Nak04, Lemma II.3.11]). Let

$$Y \xrightarrow{g} Z \xrightarrow{h} a_Y(Y) \subseteq \mathrm{Alb}(Y) \tag{4.1}$$

be the Stein factorization of a_Y.

Lemma 4.4. *If the Albanese map of Z is an algebraic fiber space, then so is that of Y.*

Proof. After possibly passing to a smooth model of Z we can assume that the Albanese map of Z is a morphism. By universality, h factors through the Albanese morphism a_Z of Z, and $a_Z \circ g$ factors through a_Y. So we have the following diagram:



By construction we have that

- $s \circ t \circ a_Z \circ g = s \circ h \circ g = s \circ a_Y = a_Z \circ g$;
- $t \circ s \circ a_Y = t \circ a_Z \circ g = h \circ g = a_Y$.

Since g is surjective, we see that $s \circ t = \mathrm{id}_{a_Z(Z)}$ and $t \circ s = \mathrm{id}_{a_Y(Y)}$. Since $a_Z(Z)$ and $a_Y(Y)$ generate $\mathrm{Alb}(Z)$ and $\mathrm{Alb}(Y)$ respectively we know that s and t are isomorphisms. Moreover since g and a_Z are both algebraic fiber spaces we are done. □

Suppose that no components of D with coefficients ≥ 1 dominates $a_Y(Y)$, or equivalently, Z. By Lemma 4.3 and [Nak04, Lemma II.3.11] we can assume Y is smooth. By [Kaw81, Theorem 13] there is an étale cover $q : Z' \to Z$ such that $Z' \cong B \times W$ where B is an abelian variety, W is birational to a smooth variety which is of general type and of maximal Albanese dimension. Let $Y' := Y \times_Z Z'$. We then do a resolution $\nu : V \to W$ and let Z'' and Y'' be the corresponding base changes. Let $\rho : Y^\sharp \to Y$ be a log resolution of $(Y'', r_*^{-1}D)$, where $r_*^{-1}D$ is the strict transform of D' on Y''. The situation is as follows.

$$\begin{array}{ccccccc}
Y^\sharp & \xrightarrow{\rho} & Y'' & \xrightarrow{g''} & Z'' = B \times V & \xrightarrow{p_V} & V \\
 & & \downarrow{\scriptstyle r} & & \downarrow{\scriptstyle s} & & \downarrow{\scriptstyle \nu} \\
 & & Y' & \xrightarrow{g'} & Z' = B \times W & \xrightarrow{p_W} & W \\
 & & \downarrow{\scriptstyle p} & & \downarrow{\scriptstyle q} & & \\
 & & Y & \xrightarrow{g} & Z & & \\
 & & \downarrow{\scriptstyle \mu} & & & & \\
 & & X & & & &
\end{array}$$

By construction V is smooth, of general type and of maximal Albanese dimension. We define $D' := p^*D$ and define $D^\sharp$ via the following inequality:

$$K_{Y^\sharp} + D^\sharp = (r \circ \rho)^*(K_{Y'} + D') + E,$$

where D' and E are effective and have no common components. Then $D^\sharp$ has snc support as well. Since p is an étale cover and $r \circ \rho$ is a birational morphism, by [Nak04, Lemma II.3.11] we have

$$0 = \kappa(K_Y + D) = \kappa(K_{Y^\sharp} + D^\sharp).$$

By Lemma 4.3, no components of $D^\sharp$ with coefficient ≥ 1 dominates Z''. We denote the horizontal part of $D^\sharp$ with respect to $g'' \circ \rho$ as $(D^\sharp)^{\mathrm{hor}}$. Then the coefficients of $(D^\sharp)^{\mathrm{hor}}$ are in $[0,1]$. By [Cam04, Theorem 4.2 and Remark 4.3] we know that

$$\kappa(K_{Y^\sharp} + (D^\sharp)^{\mathrm{hor}}) \geq \kappa((K_{Y^\sharp} + (D^\sharp)^{\mathrm{hor}})|_{F_{Y^\sharp/V}}) + \kappa(V). \tag{4.2}$$

and $\kappa(V) \geq 0$ as V is of general type.

On the other hand we have

$$\kappa(K_{F_{Y^\sharp/Z''}} + (D^\sharp)^{\mathrm{hor}}|_{F_{Y^\sharp/Z''}}) = \kappa(K_{F_{Y^\sharp/Z''}} + D^\sharp|_{F_{Y^\sharp/Z''}}) \geq 0.$$

So by Lemma 2.8, $\kappa((K_{Y^\sharp} + (D^\sharp)^{\mathrm{hor}})|_{F_{Y^\sharp/V}}) \geq 0$, and by (4.2) we have $\kappa(K_{Y^\sharp} + (D^\sharp)^{\mathrm{hor}}) \geq 0$. We also have $\kappa(K_{Y^\sharp} + D^\sharp) = 0$, which forces that $\kappa(K_{Y^\sharp} + (D^\sharp)^{\mathrm{hor}}) = 0$. Again by (4.2),

we obtain $\kappa(V) = 0$. But V is of general type by construction, so V, hence W, has to be a point. Now we consider the composition of maps

$$B \xrightarrow{q} Z \xrightarrow{h} \mathrm{Alb}(Y).$$

Since q and h are finite and $h(Z)$ generates $\mathrm{Alb}(Y)$, the composite $h \circ q$ has to be an isogeny, hence Z is birational to an abelian variety. Finally by Lemma 4.4 we are done.

For the case of the Albanese moprhism, we just do the Stein factorization as in (4.1) for a_X without taking the resolution, and the rest of the proof is the same. □

References

[BS13] M. Blickle and K. Schwede. p^{-1}-linear maps in algebra and geometry. *Commutative algebra*, pages 123–205, 2013. 8

[Cam04] F. Campana. Orbifolds, special varieties and classification theory. *Ann. Inst. Fourier (Grenoble)*, 54(3):499–630, 2004. 3, 13

[CH09] J. A. Chen and C. D. Hacon. On Ueno's conjecture K. *Math. Ann.*, 345(2):287–296, 2009. 5

[CZ15] Y. Chen and L. Zhang. The subadditivity of the Kodaira dimension for fibrations of relative dimension one in positive characteristics. *Math. Res. Lett.*, 22(3):675–696, 2015. 11

[DPS93] J-P. Demailly, T. Peternell, and M. Schneider. Kähler manifold with numerically effective Ricci class. *Compositio Math.*, 89(2):217–240, 1993. 1

[DS15] O. Das and K. Schwede. The F-different and a canonical bundle formula. *to appear in Ann. Sc. Norm. Super. Pisa*, 2015. 5

[FGI+05] B. Fantechi, L. Göttche, L. Illusie, S. L. Kleiman, N. Nitsure, and A. Vistoli. *Fundamental algebraic geometry: Grothendieck's FGA explained*. American Mathematical Society, Providence, RI, 2005. 3, 4

[Hac04] C. D. Hacon. A derived category approach to generic vanishing. *J. Reine Angew. Math.*, 575:173–187, 2004. 4

[Har77] R. Hartshorne. *Algebraic Geometry*. Springer-Verlag, New York-Heidelberg, 1977. 9

[HK10] C. D. Hacon and S. Kovács. *Classification of higher dimensional algebraic varieties*. Birkhäuser Verlag, Basel, 2010. 3

[HP13] C. D. Hacon and Z. Patakfalvi. Generic vanishing in characteristic $p > 0$ and the characterization of ordinary abelian varieties. *to appear in Amer. J. Math., arXiv:1310.2996*, 2013. 1, 2, 6, 7, 9

[HP15] C. D. Hacon and Z. Patakfalvi. On the characterization of abelian varieties in characteristic $p > 0$. *arXiv: 1602.01791*, 2015. 1, 11

[HX15] C. D. Hacon and C. Xu. On the three dimensional minimal model program in positive characteristics. *J. Amer. Math. Soc.*, 28(3):711–744, 2015. 5

[Kaw81] Y. Kawamata. Characterization of abelian varieties. *Compositio Math.*, 43(2):253–276, 1981. 1, 3, 13

[Kaw85] Y. Kawamata. Minimal models and the Kodaira dimension of algebraic fiber spaces. *J. Reine Angew. Math.*, 363:1–46, 1985. 4

[Kol13] J. Kollár. *Singularities of the minimal model program*. Cambridge University Press, Cambridge, 2013. 10

[Lan83] S. Lang. *Abelian varieties*. Springer-Verlag, New York-Berlin, 1983. 3, 4

[Moc12] S. Mochizuki. Topics in absolute anabelian geometry I: generalities. *J. Math. Sci. Univ. Tokyo*, 19(2):139–242, 2012. 8

[Muk81] S. Mukai. Duality between $D(X)$ and $D(\hat{X})$ with its application to Picard sheaves. *Nagoya Math. J.*, 81:135–175, 1981. 5

[MvdG] B. Moonen and van der Geer. *Abelian Variety*. Preliminary version, available on the authors' websites. 9

[Nak04] N. Nakayama. *Zariski-decomposition and abundance*, volume 14 of *MSJ Memoirs*. Mathematical Society of Japan, Tokyo,, 2004. 12, 13

[Pat13] Z. Patakfalvi. On subadditivity of kodaira dimesnion in positive characteristic. *to appear in J. Algebraic Geom., arXiv: 1308.5371*, 2013. 8

[Pat14] Z. Patakfalvi. Semi-positivity in positive characteristics. *Ann. Sci. École Norm Sup.*, 47(5):991–1025, 2014. 11

[PS14] M. Popa and C. Schnell. On direct images of pluricanonical bundles. *Algebra Number Theory*, 8(9):2273–2295, 2014. 3, 4

[PST14] Z. Patakfalvi, K. Schwede, and K. Tucker. Notes for the workshop on positive characteristic algebraic geometry. *arXiv: 1412.2203v1*, 2014. 5

[PSZ13] Z. Patakfalvi, K. Schwede, and W. Zhang. F-singularities in families. *arXiv: 1305.1646v1*, 2013. 11

[Rei83] M. Reid. Projective morphisms according to Kawamata. *Unpublished manuscript*, 1983. 4

[Sch09] K. Schwede. F-adjunction. *Algebra Number Theory*, 3(8):907–950, 2009. 6

[Shi15] T. Shibata. On generic vanishing for pluricanonical bundles. *arXiv: 1507.0329v4*, 2015. 3, 5

[ST11] K. Schwede and K. Tucker. A survey of test ideals. *arXiv: 1104.2000v2*, 2011. 5

[Tan15] H. Tanaka. Semiample perturbations for log canonical varieties over an F-finite field containing an infinite perfect field. *arXiv: 1503.01264v3*, 2015. 11

[Wan15a] Y. Wang. Generic vanishing and classification of irregular surfaces in positive characteristics. *to appear in Trans. Amer. Math. Soc.*, 2015. 5

[Wan15b] Y. Wang. On relative rational chain connectedness of threefolds with anti-big canonical divisors in positive characteristics. *arXiv: 1512.06189*, 2015. 2, 7

[Zha96] Q. Zhang. On projective manifolds with nef anticanonical bundles. *J. Reine Angew. Math.*, 478:57–60, 1996. 2

[Zha05] Q. Zhang. On projective varieties with nef anticanonical divisors. *Math. Ann.*, 332(3):697–703, 2005. 2, 9

Department of Mathematics, University of Utah, 155 South 1400 East, Salt Lake City, UT 84112-0090, USA

E-mail address: ywang@math.utah.edu